



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Labrador Notes.— In a box of bird-skins kindly sent me by Dr. W. T. Grenfell from Labrador in November, 1907, the following are worthy of note:

Anas platyrhynchos. MALLARD. A female or young bird taken at Nachvack in October, 1904. This duck breeds west of Hudson Bay, and is a rare transient visitor in northern Labrador.

Dafla acuta. PINTAIL. Adult male, Nachvack, June 1, 1905. This duck is a rare transient visitor in Labrador. Mr. G. M. Allen and I obtained a skin at Hopedale and saw another skin, but were able to find only five other records for Labrador. (Birds of Labrador, Proc. Boston Soc. of Nat. Hist., 1907, p. 328.)

Somateria spectabilis. KING EIDER. A curious specimen in partially albinistic plumage, shot at Battle Harbor, June 19, 1907. The whole bird is creamy white or buff, with irregular brownish patches. The belly is uniformly darker, almost vinaceous in color. The head and neck are finely lined with grayish brown streaks. The rectrices are nearly pure white, as are also the primaries and secondaries. All the feathers, especially those of the wings and tail, are much worn.

Porzana carolina. SORA. An adult taken near Harrington on the southern coast about July, 1907. This is the second record of its occurrence in Labrador. The first was of a specimen taken at Sandwich Bay in 1898. (Birds of Labrador, *loc. cit.*, p. 345.)

Falco islandus. WHITE GYRFALCON.— A very white specimen of this species taken at Nachvack, on October 13, 1905. In this connection it is interesting to note that Mr. Allen and I, who recorded this bird at Henley Harbor on the southern coast on August 2, 1906, (*loc. cit.*, p. 368) have since been told by Dr. Grenfell that a pair of these birds have bred there regularly, and that he remembers seeing them there in 1896.

Colaptes auratus luteus. NORTHERN FLICKER. A male from Sandwich Bay, just south of Hamilton Inlet, August, 1908. Mr. Allen and I concluded that this species was an "uncommon summer resident in southern half of Labrador, occasional as far as Hudson Strait" (*loc. cit.*, p. 377). The specimen is an interesting one as it appears to be somewhat darker than specimens from the New England States.

Dendroica aestiva. YELLOW WARBLER.— From Northwest River, Hamilton Inlet, taken by a cat, September 1, 1905. I hope that Mr. Oberholser will soon report on this interesting specimen.

I would also mention the following specimen kindly sent me in the autumn of 1908, by Mr. John Goleby, a Moravian brother stationed at Hopedale:

Hirundo erythrogaster. BARN SWALLOW, taken at Hopedale in July, 1908. The only previous record for Labrador is that the bird "breeds at North-west River at the head of Hamilton Inlet" (*loc. cit.*, p. 403).— CHARLES W. TOWNSEND. M. D., *Boston, Mass.*